

ABIRATERONE

Products Affected

- *abiraterone acetate*
- ZYTIGA ORAL TABLET 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant metastatic prostate cancer and used in combination with prednisone, or B.) High risk, castration-sensitive metastatic prostate cancer and used in combination with prednisone
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ACTIMMUNE

Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic granulomatous disease for use in reducing the frequency and severity of serious infections, or B.) Severe, malignant osteopetrosis (SMO)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ADEMPAS

Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form, B.) Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline), C.) Pregnancy, or D.) Patients with pulmonary hypertension associated with idiopathic interstitial pneumonia
Required Medical Information	Diagnosis of one of the following A.) Pulmonary arterial hypertension (WHO group I) and diagnosis was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.), or B.) Chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) and patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable (Female patients must be enrolled in the ADEMPAS REMS program)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AFINITOR DISPERZ

Products Affected

- AFINITOR DISPERZ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Tuberous sclerosis complex (TSC)-associated partial-onset seizures, or B.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALECENSA

Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic anaplastic lymphoma kinase (ALK) positive non-small cell lung cancer as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALKINDI

Products Affected

- ALKINDI SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of adrenocortical insufficiency and 2.) Patient requires dosages not available with other available formulations of hydrocortisone
Age Restrictions	18 years of age and younger
Prescriber Restrictions	Prescribed by or in conjunction with an endocrinologist or pediatrician
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALPHA-1 PROTEINASE INHIBITOR

Products Affected

- PROLASTIN-C INTRAVENOUS SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of alpha-1 proteinase inhibitor (alpha-1 antitrypsin) deficiency in adult patients with emphysema
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALUNBRIG

Products Affected

- ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase-positive (ALK) metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ALYQ

Products Affected

- ALYQ

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of organic nitrate or guanylate cyclase stimulators (includes intermittent use)
Required Medical Information	Diagnosis of Pulmonary arterial hypertension (PAH) confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AMBRISENTAN

Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Pregnancy, or B.) Idiopathic pulmonary fibrosis (IPF), including those with pulmonary hypertension
Required Medical Information	Diagnosis of pulmonary arterial hypertension classified as WHO Group I, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AMPHETAMINES

Products Affected

- *amphetamine-dextroamphet er*
- *dextroamphetamine sulfate er*
- *dextroamphetamine sulfate oral*
- VYVANSE

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use or use within 14 days of MAOI administration, except if prescriber is a psychiatrist with experience prescribing both MAOI and amphetamine/dextroamphetamine drugs
Required Medical Information	Diagnosis of one of the following A.) Attention deficit hyperactivity disorder (ADHD), B.) Narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), or C.) Moderate to severe binge eating disorder
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ARCALYST

Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Cryopyrin-associated periodic syndromes (CAPS), including familial cold autoinflammatory syndrome (FCAS) and Muckle-Wells Syndrome (MWS), or B.) Deficiency of interleukin-1 receptor antagonist (DIRA) and patient requires maintenance therapy for remission
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ARIKAYCE

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of pulmonary Mycobacterium avium complex (MAC) infection and used as part of a combination antibacterial regimen in treatment refractory patients
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AURYXIA

Products Affected

- AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	Iron overload syndrome (e.g. hemochromatosis)
Required Medical Information	Diagnosis of hyperphosphatemia in patients with chronic kidney disease (CKD) on dialysis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or nephrologist
Coverage Duration	12 months
Other Criteria	Ferric Citrate is NOT approvable for iron deficiency anemia per Part D law
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AUSTEDO

Products Affected

- AUSTEDO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Suicidal ideation and/or untreated or inadequately treated depression, B.) Hepatic impairment, C.) Taking MAOIs, reserpine, or tetrabenazine
Required Medical Information	Diagnosis of one of the following A.) Chorea associated with Huntington's disease (Huntington's chorea), or B.) Tardive dyskinesia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or psychiatrist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

AYVAKIT

Products Affected

- AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic gastrointestinal stromal tumor, with a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BALVERSA

Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced or metastatic urothelial carcinoma with susceptible FGFR3 or FGFR2 genetic alterations and patient has progressed during or following at least one line of prior platinum-containing chemotherapy, including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BOSENTAN

Products Affected

- *bosentan*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant cyclosporine A or glyburide therapy, or B.) Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I) and patient has New York Heart Association (NYHA) Functional Class II-IV, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with pulmonologist or cardiologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BOSULIF

Products Affected

- BOSULIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic, accelerated, or blast phase Philadelphia chromosome-positive (Ph+) chronic myelogenous leukemia (CML) with resistance or inadequate response to prior therapy, or B.) Newly diagnosed chronic phase Philadelphia chromosome-positive (Ph+) CML
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BRAFTOVI

Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) unresectable or metastatic melanoma with documented BRAF V600E or V600K mutation as detected by an FDA-approved test and used in combination with binimetinib, or B.) metastatic colorectal cancer with documented BRAF V600E mutation as detected by an FDA-approved test, patient has received prior therapy, and braftovi used in combination with cetuximab.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

BRUKINSA

Products Affected

- BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of A.) Mantle Cell Lymphoma (MCL) and patient has received at least one prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CABLIVI

Products Affected

- CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP) and used in combination with plasma exchange and immunosuppression therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CABOMETYX

Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Advanced hepatocellular carcinoma (HCC) and patient has been previously treated with sorafenib, or C.) Advanced renal cell carcinoma and used as first line treatment in combination with nivolumab
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CALQUENCE

Products Affected

- CALQUENCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) and patient has received at least 1 prior therapy, B.) Chronic lymphocytic leukemia (CLL), or C.) Small lymphocytic lymphoma (SLL)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CAPRELSA

Products Affected

- CAPRELSA

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of metastatic or unresectable locally advanced medullary thyroid cancer (MTC) AND disease is symptomatic or progressive
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CAYSTON

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (confirmed by appropriate diagnostic or genetic testing) and patient has suspected or confirmed <i>Pseudomonas aeruginosa</i> infection
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CINRYZE

Products Affected

- CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) hereditary angioedema, used in prevention of angioedema attacks, or B.) hereditary angioedema, used in prevention of acute abdominal, facial, or laryngeal attacks
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CNS STIMULANTS

Products Affected

- *armodafinil*
- *modafinil*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Obstructive sleep apnea (OSA) confirmed by sleep lab evaluation, B.) Narcolepsy confirmed by sleep lab evaluation, or C.) Shift work disorder (SWD)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COMETRIQ

Products Affected

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of progressive, metastatic medullary thyroid cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COPIKTRA

Products Affected

- COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A) chronic lymphocytic leukemia, OR B) small lymphocytic lymphoma, OR C) follicular lymphoma, AND disease is relapsed or refractory, AND patient has history of at least 2 prior therapies
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CORLANOR

Products Affected

- CORLANOR ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Decompensated acute heart failure, B.) hypotension (i.e. blood pressure less than 90/50 mmHg), C.) sick sinus syndrome or sinoatrial block or 3rd degree AV block (unless a functioning demand pacemaker is present), D.) bradycardia (i.e., resting heart rate less than 60 bpm prior to treatment), or E.) Severe hepatic impairment (Child-Pugh C)
Required Medical Information	Diagnosis of one of the following A.) stable, symptomatic chronic heart failure with left ventricular ejection fraction 35% or less, who are in sinus rhythm with resting heart rate 70 beats per minute or more and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use, or B.) stable, symptomatic heart failure due to dilated cardiomyopathy in patients who are in sinus rhythm with an elevated heart rate
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COSENTYX

Products Affected

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Ankylosing spondylitis and patient has failed or is intolerant to Humira and Enbrel, B.) Moderate to severe plaque psoriasis and patient has failed or is intolerant to Humira and Enbrel, C.) Active psoriatic arthritis and patient has failed or is intolerant to Humira and Enbrel, or D.) Non-radiographic axial spondyloarthritis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

COTELLIC

Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic malignant melanoma with BRAF V600E OR V600K mutation, and documentation of combination therapy with vemurafenib (Zelboraf)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CYSTADROPS

Products Affected

- CYSTADROPS

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Demonstrated cysteamine hypersensitivity, B.) Demonstrated penicillamine hypersensitivity
Required Medical Information	Diagnosis of cystinosis and patient has corneal cystine crystal accumulation
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

CYSTARAN

Products Affected

- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Demonstrated cysteamine hypersensitivity, B.) Demonstrated penicillamine hypersensitivity
Required Medical Information	Diagnosis of cystinosis and patient has corneal cystine crystal accumulation
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DALFAMPRIDINE

Products Affected

- *dalfampridine er*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of seizure. B.) Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute)
Required Medical Information	Diagnosis of multiple sclerosis and patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting dalfampridine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DAURISMO

Products Affected

- DAURISMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of newly diagnosed acute myeloid leukemia (AML) and used in combination with cytarabine in patients 75 years of age or older OR in patients that have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DEFERASIROX

Products Affected

- *deferasirox oral tablet*
- *deferasirox oral tablet soluble*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Creatinine clearance less than 40 mL/min, B.) Poor performance status, C.) Platelet count less than 50 x 10 ⁹ /L, D.) Advanced malignancy, E.) High-risk myelodysplastic syndrome (MDS)
Required Medical Information	Diagnosis of one of the following A.) Chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes who have liver iron concentrations of at least 5 mg Fe/g dry weight AND serum ferritin level greater than 300 mcg/L, or B.) Chronic iron overload due to blood transfusions (transfusion hemosiderosis) as evidenced by transfusion of at least 100 mL/kg packed red blood cells AND serum ferritin level greater than 1000 mcg/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DIACOMIT

Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe myoclonic epilepsy in infancy (Dravet syndrome) in patients taking clobazam
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DICLOFENAC TOPICAL

Products Affected

- *diclofenac sodium external gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Actinic keratosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DIMETHYL FUMARATE

Products Affected

- *dimethyl fumarate oral*
- *dimethyl fumarate starter pack*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DOJOLVI

Products Affected

- DOJOLVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Long-chain fatty acid oxidation disorder (LC-FAOD)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DRONABINOL

Products Affected

- dronabinol*

PA Criteria	Criteria Details
Exclusion Criteria	Sesame oil hypersensitivity
Required Medical Information	Diagnosis of one of the following A.) Anorexia associated to AIDS, or B.) Chemotherapy-induced nausea and vomiting
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

DUPIXENT

Products Affected

- DUPIXENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe atopic dermatitis and patient has trial and failure, contraindication, or intolerance to two medium to high potency topical corticosteroids (e.g., mometasone, triamcinolone, fluocinolone, betamethasone, etc), or B.) Eosinophilic phenotype or oral corticosteroid- dependent moderate to severe asthma and used as an adjunct treatment, or C.) Chronic rhinosinusitis with nasal polyposis and used as an adjunct treatment
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an allergist, dermatologist, immunologist, otolaryngologist, pulmonologist, or rheumatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SUBCUTANEOUS SOLUTION RECONSTITUTED
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, or E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENDARI

Products Affected

- ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acute sickle cell disease, or B.) Short bowel syndrome and combined with recombinant human growth hormone
Age Restrictions	5 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENSPRYNG

Products Affected

- ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Active Hepatitis B infection, or B.) Active or untreated latent tuberculosis
Required Medical Information	Diagnosis of neuromyelitis optica spectrum disorder (NMOSD) in patients who are anti-aquaporin-4 (AQP4) antibody positive
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, immunologist, or ophthalmologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ENTRESTO

Products Affected

- ENTRESTO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of angioedema related to previous ACE inhibitor or ARB therapy, B.) Concomitant use or use within 36 hours of ACE inhibitors, or C.) Concomitant use of aliskiren in patients with diabetes
Required Medical Information	Diagnosis of one of the following A.) Chronic heart failure, NYHA Class II to IV, or B.) Symptomatic heart failure with systemic left ventricular systolic dysfunction
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

EPIDIOLEX

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Lennox-Gastaut syndrome, or B.) Severe myoclonic epilepsy in infancy (Dravet syndrome), or C.) Seizures associated with tuberous sclerosis complex
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

EPOETIN THERAPY

Products Affected

- RETACRIT INJECTION SOLUTION UNIT/ML, 4000 UNIT/ML, 40000
10000 UNIT/ML, 10000 UNIT/ML(1ML), UNIT/ML
2000 UNIT/ML, 20000 UNIT/ML, 3000

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Supporting statement of diagnosis from physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ERLEADA

Products Affected

- ERLEADA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Nonmetastatic, castration-resistant prostate cancer, or B.) Metastatic, castration-sensitive prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ERLOTINIB

Products Affected

- *erlotinib hcl*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced, unresectable, or metastatic pancreatic cancer and erlotinib will be used in combination with gemcitabine, or B.) Locally advanced or metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility AND one of the following 1.) erlotinib will be used as first-line treatment, OR 2.) failure with at least one prior chemotherapy regimen, OR 3.) no evidence of disease progression after four cycles of first-line platinum-based chemotherapy and erlotinib will be used as maintenance treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ESBRIET

Products Affected

- ESBRIET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

EVEROLIMUS

Products Affected

- AFINITOR ORAL TABLET 10 MG
- *everolimus oral tablet 2.5 mg, 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Renal angiomyolipoma and tuberous sclerosis complex (TSC) not requiring immediate surgery, B.) Advanced hormone receptor-positive, HER2 negative breast cancer in postmenopausal women and taken in combination with exemestane, after failure with letrozole or anastrozole, C.) Progressive, well-differentiated, nonfunctional neuroendocrine tumors of gastrointestinal or lung origin and disease is unresectable, locally advanced, or metastatic, D.) Pancreatic progressive neuroendocrine tumors and disease is unresectable, locally advanced, or metastatic, E.) Advanced renal cell carcinoma (RCC) after failure with sunitinib or sorafenib, F.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

EVRYSDI

Products Affected

- EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of spinal muscular atrophy (SMA)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FARYDAK

Products Affected

- FARYDAK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of multiple myeloma, 2.) Medication is being used in combination with Velcade (bortezomib) and dexamethasone, 3.) Patient has received at least two prior treatment regimens, including Velcade (bortezomib) and an immunomodulatory agent [e.g., Revlimid (lenalidomide), Thalomid (thalidomide)]
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FENTANYL ORAL

Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients
Required Medical Information	Must meet all of the following 1.) Diagnosis of cancer-related breakthrough pain, 2.) Patient is currently receiving/tolerant to around-the-clock opioid therapy for persistent cancer pain, and 3.) Patient and prescriber are enrolled in the TIRF REMS Access Program
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FINTEPLA

Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI
Required Medical Information	Diagnosis of Severe myoclonic epilepsy in infancy (Dravet syndrome)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FIRAZYR

Products Affected

- FIRAZYR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary angioedema AND medication will be used for the treatment of acute attacks.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

FIRDAPSE

Products Affected

- FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	History of seizures
Required Medical Information	Diagnosis of Lambert-Eaton myasthenic syndrome
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GALAFOLD

Products Affected

- GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Fabry disease with an amenable galactosidase alpha gene (GLA) mutation
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GATTEX

Products Affected

- GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of short bowel syndrome and patient is dependent on parenteral support
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GAVRETO

Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC) as detected by an FDA approved test, B.) Advanced or metastatic RET-mutant medullary thyroid cancer and patient requires systemic therapy, or C.) Advanced or metastatic RET fusion-positive thyroid and patient requires systemic therapy and is radioactive iodine-refractory, when radioactive iodine is appropriate
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GILENYA

Products Affected

- GILENYA ORAL CAPSULE 0.5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure, B.) History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker, C.) Baseline QTC interval greater than or equal to 500 milliseconds, D.) Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol)
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	10 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GILOTRIF

Products Affected

- GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have nonresistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test, or B.) Metastatic squamous NSCLC with progression after platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GLATIRAMER

Products Affected

- COPAXONE SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- *glatiramer acetate*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GOCOVRI

Products Affected

- GOCOVRI

PA Criteria	Criteria Details
Exclusion Criteria	Patients with end-stage renal disease (ESRD, CrCl below 15 ml/min/m ²)
Required Medical Information	Diagnosis of one of the following A.) Parkinson disease and patient is experiencing dyskinesia, receiving levodopa based therapy, and has documented trial and failure to amantadine immediate release, B.) Extrapyrimalidal disease and has documented trial and failure to amantadine immediate release, or C.) Parkinson disease and patient is experiencing "off" episodes, receiving levodopa/carbidopa based therapy, and has documented trial and failure to amantadine immediate release
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

GROWTH HORMONE

Products Affected

- OMNITROPE SUBCUTANEOUS SOLUTION CARTRIDGE
- OMNITROPE SUBCUTANEOUS SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Use for growth promotion in pediatric patients with closed epiphyses, B.) Acute critical illness caused by complications following open-heart or abdominal surgery, multiple accidental trauma, or acute respiratory failure, C.) Active malignancy, D.) Active proliferative or severe nonproliferative diabetic retinopathy, E.) Prader-Willi Syndrome in patients who are severely obese, have a history of upper airway obstruction or sleep apnea, or have severe respiratory impairment
Required Medical Information	Diagnosis of pediatric indication: A.) GHD and bone age at least 1 year or 2 standard deviations (SD) delayed compared with chronological age and 2 stim tests with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SDS below mean if CNS pathology, h/o irradiation, or proven genetic cause, B.) SGA and birth weight or length 2 or more SDS below mean for gestational age and fails to manifest catch up growth by age 2 (height 2 or more SDS below mean for age and gender), C.) CRI and nutritional status has been optimized, metabolic abnormalities have been corrected, and patient has not had renal transplant D.) SHOX deficiency or Noonan syndrome E.) PWS confirmed by genetic testing, F.) Turner Syndrome confirmed by chromosome analysis. For GHD, CRI, SHOX deficiency, Noonan syndrome, and PWS one of the following height more than 3 SDS below mean for age and gender, or height more than 2 SDS below mean with GV more than 1 SDS below mean, or GV over 1 year 2 SDS below mean. OR Diagnosis of an adult indication: A.) childhood- or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone [preferred], glucagon, arginine), B.) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C.) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D.) GHD with irreversible

PA Criteria	Criteria Details
	hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region AND a subnormal IGF-1 (after at least 1 month off GH therapy) AND Objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or cardiovascular complications AND Completed linear growth (GV less than 2 cm/year) AND GH has been discontinued for at least 1 month (if previously receiving GH)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an Endocrinologist or Nephrologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HEMADY

Products Affected

- HEMADY

PA Criteria	Criteria Details
Exclusion Criteria	Systemic fungal infections
Required Medical Information	Diagnosis of multiple myeloma (MM), used in combination with other anti-myeloma drugs, and treatment regimen cannot be supported by lower strengths of oral dexamethasone
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HEPATITIS C

Products Affected

- MAVYRET
- *sofosbuvir-velpatasvir*
- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of HCV genotype, subtype and quantitative HCV RNA (viral load) testing any time prior to therapy. Must document the following within 12 weeks of starting therapy: CBC, INR, hepatic function panel, and GFR. Must document cirrhosis status, prior treatment history (if any), and planned duration of treatment. All genotypes will require trial/failure, contraindication to, or intolerance to Mavyret or Sofosbuvir-Velpatasvir prior to the approval of Vosevi.
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HETLIOZ

Products Affected

- HETLIOZ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Non-24-hour-sleep-wake disorder (Non-24), or B.) Nighttime sleep disturbances associated with Smith-Magenis Syndrome (SMS)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HRM - SKELETAL MUSCLE RELAXANTS

Products Affected

- chlorzoxazone oral tablet 375 mg, 500 mg, 750 mg
- cyclobenzaprine hcl oral
- methocarbamol oral
- orphenadrine citrate er

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	For patients greater than or equal to 65 years, coverage determination is approved for FDA-approved indications not otherwise excluded from Part D.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

HUMIRA

Products Affected

- HUMIRA PEDIATRIC CROHNS START SUBCUTANEOUS PREFILLED SYRINGE KIT 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML
- HUMIRA PEN SUBCUTANEOUS PEN-INJECTOR KIT
- HUMIRA PEN-CD/UC/HS STARTER
- HUMIRA PEN-PS/UV/ADOL HS START SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML
- HUMIRA PEN-PSOR/UEIT STARTER
- HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy, G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants (e.g. corticosteroids, azathioprine), H.) Non-infectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment



Local.
Reliable.
Accessible.

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IBRANCE

Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with fulvestrant and disease has progressed following endocrine therapy, or B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with an aromatase inhibitor in postmenopausal women or men
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ICLUSIG

Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated, B.) Chronic phase, chronic myeloid leukemia (CML) in adult patients with resistance or intolerance to at least two prior kinase inhibitors, or C.) Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IDHIFA

Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase 2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IMBRUVICA

Products Affected

- IMBRUVICA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) and patient has received at least one prior therapy, B.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL), C.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion, D.) Waldenstrom's macroglobulinemia (WM), E.) Marginal zone lymphoma (MZL) and patient requires systemic therapy and has received at least one prior anti-CD20-based therapy, or F.) Chronic graft vs host disease (cGVHD) after failure of a least one first-line corticosteroid therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INBRIJA

Products Affected

- INBRIJA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concurrent use with nonselective monoamine oxidase inhibitors (MAOIs) (e.g. phenelzine and tranylcypromine), B.) Recent use (within 2 weeks) with a nonselective MAOI
Required Medical Information	Must meet all of the following A.) Diagnosis of Parkinson's disease, B.) Concurrent therapy with carbidopa/levodopa, C.) Patient has tried and failed or has contraindication to one generic formulary alternative
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INCRELEX

Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) active or suspected malignancy, B.) use for growth promotion in patients with closed epiphyses, C.) Intravenous administration
Required Medical Information	Prescribed for treatment of growth failure in pediatric patient AND patient has diagnosis of one of the following A.) Severe primary insulin-like growth factor-1 (IGF-1) deficiency, or B.) Growth hormone (GH) gene deletion and patient has developed neutralizing antibodies to GH
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INQOVI

Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of myelodysplastic syndromes (MDS), including previously treated and untreated, de novo and secondary MDS with the following French-American-British subtypes (refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, and chronic myelomonocytic leukemia [CMML]) and intermediate-1, intermediate-2, and high-risk International Prognostic Scoring System groups
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INREBIC

Products Affected

- INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate-2 or high-risk primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis (MF).
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

INTRAROSA

Products Affected

- INTRAROSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, or B.) Known or suspected estrogen-dependent neoplasia
Required Medical Information	Diagnosis of one of the following A.) moderate to severe dyspareunia due to menopause, or B.) atrophic vaginitis due to menopause
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 3 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IRESSA

Products Affected

- IRESSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) and must meet both of the following 1.) tumor has epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility, AND 2.) Used as first-line treatment
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ISTURISA

Products Affected

- ISTURISA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing's disease in patients for whom pituitary surgery is not an option or has not been curative
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ITRACONAZOLE

Products Affected

- *itraconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.)
Required Medical Information	Diagnosis of one of the following A.) Systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), or B.) Onychomycosis confirmed by one of the following positive potassium hydroxide (KOH) preparation, fungal culture, or nail biopsy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ITRACONAZOLE SOLN

Products Affected

- *itraconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.)
Required Medical Information	Diagnosis of candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

IVIG

Products Affected

- GAMMAGARD
- GAMMAGARD S/D LESS IGA
- GAMUNEX-C
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 2 GM/20ML
- PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) IgA deficiency with antibody formation and a history of hypersensitivity, or B.) History of anaphylaxis or severe systemic reaction to human immune globulin
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

JAKAFI

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis, B.) Polycythemia vera AND patient has had an inadequate response to or is intolerant of hydroxyurea, OR C.) Acute graft versus host disease AND disease is refractory to steroid therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

JUXTAPID

Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests, B.) Pregnancy, or C.) Concomitant use with strong or moderate CYP3A4 inhibitors
Required Medical Information	Diagnosis of HoFH as confirmed by one of the following A.) Genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (LDLRAP1 or ARH), or B.) Both of the following 1.) Either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL, and 2.) Either xanthoma before 10 years of age or evidence of heterozygous FH in both parents
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KALYDECO

Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KESIMPTA

Products Affected

- KESIMPTA

PA Criteria	Criteria Details
Exclusion Criteria	Active Hepatitis B infection
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KISQALI

Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Hormone receptor (HR)-positive, HER-2 negative advanced or metastatic breast cancer in pre/perimenopausal or postmenopausal women and used in combination with an aromatase inhibitor, or B.) Hormone receptor (HR)-positive, HER-2 negative advanced or metastatic breast cancer in postmenopausal women and used in combination with fulvestrant
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KISQALI FEMARA

Products Affected

- KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KISQALI FEMARA(200 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR)-positive, HER-2 negative advanced or metastatic breast cancer in pre/perimenopausal or postmenopausal women
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KORLYM

Products Affected

- KORLYM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) pregnancy, B.) coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, C.) concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, D.) history of unexplained vaginal bleeding, E.) endometrial hyperplasia with atypia or endometrial carcinoma
Required Medical Information	Diagnosis of endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and both of the following 1.) Used to control hyperglycemia secondary to hypercortisolism, AND 2.) Patient has failed or is not a candidate for surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KOSELUGO

Products Affected

- KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) in a patient who has symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	2 years of age to 17 years of age
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

KUVAN

Products Affected

- KUVAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 2 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LENVIMA

Products Affected

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer, B.) Advanced renal cell carcinoma, in combination with everolimus, following one prior anti-angiogenic therapy, C.) Unresectable hepatocellular carcinoma, first-line therapy, D.) Advanced endometrial carcinoma that is not microsatellite instability-high or mismatch repair deficient, in combination with pembrolizumab, when disease has progressed following prior systemic therapy AND patient is not a candidate for curative surgery or radiation
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LEUKINE

Products Affected

- LEUKINE INJECTION SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Patient has undergone allogeneic or autologous bone marrow transplant (BMT) and engraftment is delayed or failed, B.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, C.) Medication will be used for myeloid reconstitution after an autologous or allogeneic BMT, D.) Patient has acute myeloid leukemia and administration will be after completion of induction chemotherapy, E.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS) or F.) Autologous peripheral blood stem cell transplant, Following myeloablative chemotherapy.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LEUPROLIDE

Products Affected

- *leuprolide acetate injection*
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic prostate cancer and patient has failed or is intolerant to Eligard (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), B.) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only), C.) Anemia due to uterine leiomyomata (Fibroids) (3.75 mg 1-month & 11.25 mg 3-month depots only) and patient is preoperative, or D.) Central precocious puberty (idiopathic or neurogenic) in children
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LIDOCAINE PATCH

Products Affected

- *lidocaine external patch 5 %*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Pain associated with diabetic neuropathy, B.) Pain associated with cancer-related neuropathy, C.) Post-herpetic neuralgia, D.) Back pain, or E.) Osteoarthritis of the knee or hip
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LONSURF

Products Affected

- LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy, or B.) Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least 2 prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LORBRENA

Products Affected

- LORBRENA

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A4 inducers
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) and one of the following 1.) Disease has progressed on alectinib as the first ALK inhibitor therapy for metastatic disease, OR 2.) Disease has progressed on ceritinib as the first ALK inhibitor therapy for metastatic disease, OR 3.) Disease has progressed on crizotinib AND at least 1 other ALK inhibitor for metastatic disease
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

LYNPARZA

Products Affected

- LYNPARZA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) HER2-negative, deleterious or suspected deleterious germline BRCA mutated metastatic breast cancer AND patient has been previously treated with chemotherapy in neoadjuvant, adjuvant, or metastatic setting, B.) Advanced ovarian cancer with known or suspected BRCA mutation as detected by an FDA-approved test AND patient has trial and failure, contraindication, or intolerance to 3 or more prior lines of chemotherapy, C.) Recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer AND used for maintenance treatment in patients who are in complete or partial response to platinum-based chemotherapy (e.g. cisplatin, carboplatin), D.) Deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients with complete or partial response to first-line platinum-based chemotherapy, E.) Deleterious or suspected deleterious germline BRCA-mutated metastatic pancreatic adenocarcinoma and disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen, F.) Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA-mutation, and/or genomic instability. Used in combination with bevacizumab for maintenance treatment., or G.) Deleterious or suspected deleterious germline or somatic homologous recombination repair gene mutated metastatic castration-resistant prostate cancer in patients who have progressed following prior treatment with enzalutamide or abiraterone.
Age Restrictions	None

PA Criteria	Criteria Details
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MAYZENT

Products Affected

- MAYZENT

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) CYP2C9*3/*3 genotype, B.) In the last 6 months experienced myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, Class III-IV heart failure, or C.) Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome, unless patient has a functioning pacemaker
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis, including clinically isolated syndrome, relapsing-remitting disease, or active secondary progressive disease and the following A.) Patients with relapsing forms of multiple sclerosis have history of/or contraindication to Avonex, Betaseron, Copaxone, Gilenya, or Tecfidera
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MEKINIST

Products Affected

- MEKINIST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and used in combination with dabrafenib and no locoregional treatment options, B.) Malignant melanoma with lymph node involvement and following complete resection with BRAF V600E or V600K mutations and used in combination with dabrafenib, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutations and used in combination with dabrafenib or as monotherapy , or D.) Metastatic non-small cell lung cancer, with BRAF V600E mutation, in combination with dabrafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MEKTOVI

Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable or metastatic malignant melanoma with documented BRAF V600E or V600K mutation as detected by an FDA approved test AND used in combination with encorafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

METHYLPHENIDATES

Products Affected

- *methylphenidate hcl er*
- *methylphenidate hcl er (cd)*
- *methylphenidate hcl er (la)*
- *methylphenidate hcl er (xr)*
- *methylphenidate hcl oral solution*
- *methylphenidate hcl oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Symptoms of marked anxiety, tension, or agitation, B.) Glaucoma, C.) Family history/diagnosis of Tourette's syndrome or presence of motor tics, D.) Concurrent use with MAOIs
Required Medical Information	Diagnosis of one of the following A.) Attention deficit hyperactivity disorder (ADHD), or B.) Narcolepsy as confirmed by a sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MIGLUSTAT

Products Affected

- *miglustat*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

MS INTERFERONS

Products Affected

- AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT
- AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NATPARA

Products Affected

- NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hypoparathyroidism and used to control hypocalcemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NERLYNX

Products Affected

- NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Early stage HER2-positive breast cancer and used following adjuvant trastuzumab therapy, or B.) Advanced or metastatic HER2-positive breast cancer, used in combination with capecitabine, AND patient has received 2 or more prior anti-HER2-based regimens in the metastatic setting
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NEXAVAR

Products Affected

- NEXAVAR

PA Criteria	Criteria Details
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma that is refractory to radioactive iodine treatment, or C.) Unresectable hepatocellular carcinoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NINLARO

Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma, used in combination with lenalidomide and dexamethasone, AND patient has history of at least 1 prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NORTHERA

Products Affected

- NORTHERA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (e.g., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUBEQA

Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of non-metastatic, castration-resistant prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUCALA

Products Affected

- NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Severe asthma with eosinophilic phenotype, B.) Eosinophilic granulomatosis with polyangiitis (EGPA), or C.) Hypereosinophilic syndrome lasting at least 6 months without an identifiable non-hematologic secondary cause
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, immunologist, pulmonologist, or rheumatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUEDEXTA

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of prolonged QT interval, congenital long QT syndrome or Torsades de pointes, B.) Heart failure, C.) Complete AV block without an implanted pacemaker or high risk of complete AV block, D.) Concomitant use with quinidine, quinine, mefloquine, or drugs that prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), E.) Concomitant use with MAOIs or within 14 days of MAOI therapy
Required Medical Information	Diagnosis of pseudobulbar affect
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

NUPLAZID

Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hallucinations and delusions associated with Parkinson disease psychosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OCTREOTIDE

Products Affected

- *octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acromegaly and patient has inadequate response to or is ineligible for surgery, radiation, or bromocriptine mesylate, or B.) Metastatic carcinoid syndrome, or C.) Vasoactive intestinal peptide-secreting tumors (VIPomas) with associated diarrhea
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ODOMZO

Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of locally advanced basal cell carcinoma of the skin and one of the following A.) Cancer has recurred following surgery or radiation therapy, B.) Patient is not a candidate for surgery or radiation therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OFEV

Products Affected

- OFEV

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Idiopathic pulmonary fibrosis (IPF), B.) Systemic sclerosis-associated interstitial lung disease (ILD), or C.) Chronic fibrosing interstitial lung disease with a progressive phenotype
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ONUREG

Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute myeloid leukemia (AML) used in maintenance treatment for adult patients who achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OPSUMIT

Products Affected

- OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ORGOVYX

Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ORILISSA

Products Affected

- ORILISSA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Pregnancy, B.) Known osteoporosis, C.) Severe hepatic impairment, D.) Concurrent use of strong organic anion transporting polypeptide (OATP) 1B1 inhibitors
Required Medical Information	Diagnosis of moderate to severe pain associated with endometriosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ORKAMBI

Products Affected

- ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OSPHENA

Products Affected

- OSPHENA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Undiagnosed abnormal genital bleeding, B.) Known or suspected estrogen-dependent neoplasia, C.) Active deep vein thrombosis (DVT), pulmonary embolism (PE), or a history of these conditions, D.) Active arterial thromboembolic disease (eg. stroke, myocardial infarction) or a history of these conditions, or E.) Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause, or B.) Moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

OXANDROLONE

Products Affected

- *oxandrolone oral*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Known or suspected carcinoma of the prostate or breast in males, B.) Carcinoma of the breast in females with hypercalcemia, C.) Pregnancy, D.) Nephrosis or nephrotic phase of nephritis, E.) Hypercalcemia
Required Medical Information	Diagnosis of one of the following A.) Bone pain associated with osteoporosis, B.) Protein catabolism associated with chronic corticosteroid administration, or C.) Used as adjunctive therapy to promote weight gain after weight loss associated with one of the following 1.) Extensive surgery, 2.) Chronic infections, 3.) Severe trauma, or 4.) Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PEGYLATED INTERFERON

Products Affected

- PEGASYS SUBCUTANEOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon, B.) Uncontrolled depression
Required Medical Information	Diagnosis of one of the following A.) Chronic hepatitis B infection, or B.) Chronic hepatitis C and required criteria will be applied consistent with current AASLD-IDSA guidance with compensated liver disease
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	HBV: 12 months, HCV: based on current AASLD guidelines
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PEMAZYRE

Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with confirmed fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, gastroenterologist, or hepatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PIQRAY

Products Affected

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR) positive, HER2-negative, PIK3CA-mutated, advanced or metastatic breast cancer AND must meet all of the following 1.) Used in combination with fulvestrant, AND 2.) Disease has progressed on or after an endocrine-based regimen, AND 3.) Patient is a male OR postmenopausal female
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

POMALYST

Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) AIDS-related Kaposi sarcoma and patient has failure on highly active antiretroviral therapy (HAART), B.) Kaposi sarcoma in HIV-negative adults, or C.) Multiple myeloma and in combination with dexamethasone in adults who have received at least 2 prior therapies (including lenalidomide and a proteasome inhibitor) and have demonstrate disease progression on or within 60 days of completion of the last therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PROMACTA

Products Affected

- PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic thrombocytopenic purpura (ITP), B.) Chronic hepatitis C infection associated thrombocytopenia, or C.) Severe aplastic anemia with insufficient response to immunosuppressive therapy or in combination with standard immunosuppressive therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

QINLOCK

Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced gastrointestinal stromal tumor (GIST) and patient has received prior treatment with 3 or more kinase inhibitors, including imatinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

REGRANEX

Products Affected

- REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	Known neoplasm at the site of application
Required Medical Information	Diagnosis of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

REPATHA

Products Affected

- REPATHA
- REPATHA SURECLICK
- REPATHA PUSHTRONEX SYSTEM

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) primary hyperlipidemia including heterozygous familial hypercholesterolemia (HeFH), B.) homozygous familial hypercholesterolemia, C.) established cardiovascular disease and myocardial infarction prophylaxis, stroke prophylaxis, or coronary revascularization prophylaxis is required, or D.) clinical atherosclerotic cardiovascular disease (CVD) as defined as one of the following 1.) acute coronary syndrome, 2.) history of myocardial infarction, 3.) stable/unstable angina, 4.) coronary or other arterial revascularization, 5.) stroke, 6.) transient ischemic stroke (TIA), or 7.) peripheral arterial disease presumed to be atherosclerotic region
Age Restrictions	13 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 2 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RETEVMO

Products Affected

- RETEVMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in patients who require systemic therapy, B.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC), or C.) Advanced or metastatic RET fusion-positive thyroid cancer in patients who require systemic therapy and are refractory to radioactive iodine, if appropriate
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

REVLIMID

Products Affected

- REVLIMID

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality or without additional cytogenetic abnormalities, D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ROZLYTREK

Products Affected

- ROZLYTREK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) ROS1-positive metastatic non-small cell lung cancer (NSCLC), or B.) Solid tumors that have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, are metastatic or where surgical resection is likely to result in severe morbidity, and have either progressed following treatment or have no satisfactory alternative therapy
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RUBRACA

Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Epithelial ovarian, fallopian tube, or primary peritoneal cancer with deleterious BRCA mutation (germline and/or somatic) as detected by an FDA-approved test and patient has been treated with 2 or more prior lines of chemotherapy, B.) Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, used as maintenance treatment, and patient is in complete or partial response to platinum-based chemotherapy, or C.) Deleterious BRCA mutation (germline and/or somatic)-associated metastatic castration-resistant prostate cancer and patient has been treated with androgen receptor-directed therapy and a taxane-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

RYDAPT

Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation therapy, or B.) systemic mastocytosis or mast cell leukemia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SAMSCA

Products Affected

- SAMSCA
- tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Diagnosis of Autosomal Dominant Polycystic Kidney Disease (ADPKD), B.) Urgent need to raise serum sodium acutely, C.) Inability to sense or appropriately respond to thirst, D.) Hypovolemic hyponatremia, E.) Concomitant use of strong CYP 3A Inhibitors (eg. clarithromycin, ketoconazole, ritonavir), F.) Anuria
Required Medical Information	Diagnosis of clinically significant hypervolemic or euvolemic hyponatremia (serum sodium less than 125 mEq/L or less marks hyponatremia that is symptomatic and has resisted correction with fluid restriction), including in patients with heart failure and syndrome of inappropriate antidiuretic hormone (SIADH)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SAPROPTERIN

Products Affected

- *sapropterin dihydrochloride*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 2 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SIGNIFOR

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing disease and patient has had inadequate response to or is not a candidate for surgery. For renewal: Documentation of a clinically meaningful reduction in 24-hour urinary free cortisol (UFC) levels or improvement in signs or symptoms of the disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 6 months, Reauthorization: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SILDENAFIL

Products Affected

- *sildenafil citrate oral tablet 20 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Nitrate therapy, including intermittent use
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SOMATULINE DEPOT

Products Affected

- SOMATULINE DEPOT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acromegaly and patient is not a candidate for surgery/radiotherapy or has had an inadequate response, B.) Carcinoid syndrome, or C.) Unresectable, well or moderately differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SOMAVERT

Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly and patient has had an inadequate response to or is ineligible for surgery or radiation therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SPRYCEL

Products Affected

- SPRYCEL

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, B.) Chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy, C.) Ph+ acute lymphoblastic leukemia (ALL) with resistance or intolerance to prior therapy, D.) Newly diagnosed Ph+ ALL in combination with chemotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

STELARA

Products Affected

- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severely active Crohn disease and patient has trial and failure or intolerance or contraindication to Humira, B.) Moderate to severe plaque psoriasis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, C.) Active psoriatic arthritis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, or D.) Moderate to severe active ulcerative colitis and patient has trial and failure or intolerance or contraindication to Humira.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

STIVARGA

Products Affected

- STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer in patients previously treated with fluoropyrimidine, oxaliplatin, and irinotecan containing chemotherapy, anti-VEGF therapy, and if RAS wild type, anti-EGFR therapy, B.) Liver carcinoma in patients previously treated with sorafenib, or C.) Locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with imatinib and sunitinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SUNOSI

Products Affected

- SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI
Required Medical Information	Diagnosis of one of the following A.) narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to modafinil or armodafinil, or B.) obstructive sleep apnea (OSA) with excessive daytime drowsiness and has trial of/or contraindication to modafinil or armodafinil
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SUTENT

Products Affected

- SUTENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Gastrointestinal stromal tumor after disease progression on or intolerance to imatinib, B.) Pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease, C.) Advanced renal cell carcinoma, or D.) Renal cell carcinoma and used as adjuvant therapy following nephrectomy in patients who are at high risk for recurrence
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SYMDEKO

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and must meet one of the following 1.) Patient is homozygous for the F508del mutation, or 2.) Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	Initial: 6 months, Renewal: 12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

SYMLIN

Products Affected

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Confirmed diagnosis of gastroparesis, B.) Hypoglycemia unawareness
Required Medical Information	Diagnosis of type 1 or type 2 diabetes mellitus and patient uses mealtime insulin therapy and has failed to achieve desired glucose control
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TABRECTA

Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAFINLAR

Products Affected

- TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid carcinoma with BRAF V600E mutation, in combination with trametinib and no satisfactory locoregional treatment options, B.) Metastatic non-small cell lung cancer with BRAF V600E mutation, in combination with trametinib OR in patients previously treated as monotherapy, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAGRISO

Products Affected

- TAGRISO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R mutation and used as first line therapy, B.) Metastatic non-small cell lung cancer with T790M EGFR mutation (as confirmed by an FDA-approved test) AND whose disease has progressed on or after EGFR tyrosine kinase inhibitor therapy, or C.) Non-small cell lung cancer (NSCLC) with tumor epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations (as confirmed by an FDA-approved test) AND patient requires adjuvant therapy after tumor resection
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAKHZYRO

Products Affected

- TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary angioedema, used in prevention of angioedema attacks AND patient has trial of, or contraindication to Firazyr
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TALZENNA

Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious or suspected deleterious germline breast cancer susceptibility gene (BRCA)-mutated (gBRCAm), human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TARGRETIN GEL

Products Affected

- TARGRETIN EXTERNAL

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL Stage 1A/1B) and patient had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) indicated for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TASIGNA

Products Affected

- TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: A.) Long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (CML), B.) Chronic phase or accelerated phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior therapy that included imatinib, or C.) Chronic phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior tyrosine-kinase inhibitor therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TAZVERIK

Products Affected

- TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic or locally advanced epithelioid sarcoma in patients not eligible for complete resection, B.) Relapsed or refractory follicular lymphoma in patients whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies, or C.) Relapsed or refractory follicular lymphoma in patients who have no satisfactory alternative treatment options
Age Restrictions	16 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TECFIDERA

Products Affected

- TECFIDERA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TEGSEDI

Products Affected

- TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Platelet count less than 100,000 per microliter, B.) Urinary protein to creatinine ratio (UPCR) of 1000 mg/g or higher
Required Medical Information	Diagnosis of Polyneuropathy of hereditary transthyretin-mediated amyloidosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TERIPARATIDE

Products Affected

- *teriparatide (recombinant)*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Patient has previous trial and failure, contraindication, or intolerance to a bisphosphonate AND diagnosis of one of the following A.) osteoporosis in postmenopausal female patient with high risk for fracture and patient has history of or contraindication to Tymlos, B.) primary or hypogonadal osteoporosis in male patient with high risk for fracture, or C.) osteoporosis due to associated sustained systemic glucocorticoid therapy in patient with high risk for fracture
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: 12 months (Maximum 24 month treatment per patient lifetime)
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TETRABENAZINE

Products Affected

- *tetrabenazine*

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Actively suicidal, B.) Untreated or inadequately treated depression, C.) Impaired hepatic function, D.) Concomitant use of monoamine oxidase inhibitors, E.) Concomitant use of reserpine or within 20 days of discontinuing reserpine
Required Medical Information	Diagnosis of chorea associated with Huntington's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

THALOMID

Products Affected

- THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma that is newly diagnosed, or B.) Erythema nodosum leprosum (ENL)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or infectious disease specialist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TIBSOVO

Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory acute myeloid leukemia with a susceptible isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test), or B.) Newly diagnosed acute myeloid leukemia with susceptible isocitrate dehydrogenase-1 mutation AND meets one of the following 1.) Patient is 75 years of age or older, OR 2.) Patient has comorbidities that preclude intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRIENTINE

Products Affected

- CLOVIQUE
- trientine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease in patients that are intolerant to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TRIKAFTA

Products Affected

- TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and patient has at least 1 F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene verified by an FDA-cleared CF mutation test
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TUKYSA

Products Affected

- TUKYSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced unresectable or metastatic HER2-positive breast cancer (including brain metastases) in patients who have received one or more prior anti-HER2-based regimens in the metastatic setting and drug is being used in combination with trastuzumab and capecitabine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TURALIO

Products Affected

- TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

TYMLOS

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of postmenopausal osteoporosis and one of the following A.) osteoporotic fracture or multiple risk factors for fracture, or B.) previous trial of/or contraindication to bisphosphonate
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 12 months, Renewal: 12 months (Maximum 24 month treatment per patient lifetime)
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

UPTRAVI

Products Affected

- UPTRAVI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP2C8 inhibitors (e.g., gemfibrozil)
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VENCLEXTA

Products Affected

- VENCLEXTA
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A inhibitor during the initial and titration phase in patients with CLL or SLL
Required Medical Information	Diagnosis of one of the following A.) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL), or B.) Newly-diagnosed acute myeloid leukemia (AML) and used in combination with azacitidine, decitabine or low-dose cytarabine in patients 75 years or older or who have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VERZENIO

Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic, HER2-negative, hormone receptor-positive breast cancer AND one of the following: A.) For postmenopausal women must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy and patient has trial and failure or contraindication to Ibrance or Kisqali, B.) For premenopausal or perimenopausal women must be used in combination with fulvestrant for the treatment of disease progression following endocrine therapy and patient has trial and failure or contraindication to Ibrance, C.) Used as monotherapy for treatment of disease progression following endocrine therapy and patient has already received at least one prior chemotherapy regimen of Ibrance or Kisqali, D.) For postmenopausal women used as initial endocrine-based treatment in combination with an aromatase inhibitor and patient has trial and failure or contraindication to Kisqali or Ibrance, E.) For premenopausal or perimenopausal women used as initial endocrine-based treatment in combination with an aromatase inhibitor and patient has trial and failure or contraindication to Kisqali
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off-Label Uses	N/A

VITRAKVI

Products Affected

- VITRAKVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or surgically unresectable neurotrophic receptor tyrosine kinase (NTRK) gene fusion positive solid tumors and used in patients with unsatisfactory alternative treatments or who have progressed following treatment
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VIZIMPRO

Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer with confirmed epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VORICONAZOLE

Products Affected

- *voriconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Invasive aspergillosis, B.) Candidemia, C.) Esophageal Candidiasis, D.) Invasive candidiasis of the skin and abdomen, kidney, bladder wall, and wounds, or E.) Serious fungal infection due to <i>Scedosporium apiospermum</i> or <i>Fusarium</i> species
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	6 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VOTRIENT

Products Affected

- VOTRIENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, or B.) Advanced soft tissue sarcoma and patient received at least one prior chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

VYNDAMAX

Products Affected

- VYNDAMAX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of wild type or hereditary transthyretin related familial amyloid cardiomyopathy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XALKORI

Products Affected

- XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive as detected by an FDA-approved test, or B.) Relapsed or refractory systemic anaplastic large cell lymphoma that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XELJANZ

Products Affected

- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) Moderate to severe rheumatoid arthritis (RA) and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, B.) Active psoriatic arthritis and patient has trial and failure or intolerance or contraindication to Humira and Enbrel, or C.) Moderate to severe ulcerative colitis (UC) and patient has trial and failure or intolerance or contraindication to Humira.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XGEVA

Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia (calcium less than 8.0 mg/dL)
Required Medical Information	Diagnosis of one of the following A.) Bone metastases from a solid tumor and used for the prevention of skeletal related events, B.) Multiple myeloma and used for the prevention of skeletal related events, C.) Hypercalcemia of malignancy refractory to bisphosphonate therapy, or D.) Giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XOLAIR

Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic urticaria in patients who remain symptomatic despite H1 antihistamine therapy, B.) Moderate to severe persistent asthma in patients with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms are inadequately controlled with inhaled corticosteroids, or C.) Nasal polyps in patients with inadequate response to nasal corticosteroids
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, dermatologist, immunologist, otolaryngologist, or pulmonologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XOSPATA

Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XPOVIO

Products Affected

- XPOVIO (100 MG ONCE WEEKLY)
- XPOVIO (40 MG ONCE WEEKLY)
- XPOVIO (40 MG TWICE WEEKLY)
- XPOVIO (60 MG ONCE WEEKLY)
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY)
- XPOVIO (80 MG TWICE WEEKLY)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory multiple myeloma being used in combination with dexamethasone in a patient who has received at least 4 prior therapies and is refractory to at least 2 proteasome inhibitors, at least 2 immunomodulatory agents, and an anti-CD38 monoclonal antibody, B.) Relapsed or refractory diffuse large B-cell lymphoma (DLBCL, including from follicular lymphoma) in a patient who has received at least 2 lines of systemic therapy, or C.) Multiple myeloma being used in combination with bortezomib and dexamethasone in a patient who has received at least 1 prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XTANDI

Products Affected

- XTANDI ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant prostate cancer, or B.) Metastatic, castration-sensitive prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XURIDEN

Products Affected

- XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary orotic aciduria
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XYREM

Products Affected

- XYREM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency
Required Medical Information	Diagnosis of one of the following A.) narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to modafinil or armodafinil, or B.) cataplexy and narcolepsy
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

XYWAV

Products Affected

- XYWAV

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency
Required Medical Information	Diagnosis of one of the following A.) Narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to modafinil or armodafinil, or B.) Cataplexy and narcolepsy
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

YONSA

Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of metastatic, castration-resistant prostate cancer and use in combination with methylprednisolone
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZARXIO

Products Affected

- ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chemotherapy induced febrile neutropenia (prophylaxis), B.) Severe chronic neutropenia, C.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, or D.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZEJULA

Products Affected

- ZEJULA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) advanced or recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer and used for maintenance therapy in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin), or B.) advanced ovarian, fallopian tube, or primary peritoneal cancer and patient has been treated with 3 or more prior chemotherapy regimens, and cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA mutation, or genomic instability, and disease has progressed more than 6 months after response to the last platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or gynecologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZIEXTENZO

Products Affected

- ZIEXTENZO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chemotherapy induced febrile neutropenia (prophylaxis)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

ZYKADIA

Products Affected

- ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A

PART B VERSUS PART D

Products Affected

- ABELCET INTRAVENOUS SUSPENSION 5 MG/ML
- *acetylcysteine inhalation solution 10 %, 20 %*
- *acyclovir sodium intravenous solution 50 mg/ml*
- *albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml*
- AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG
- AMINOSYN-PF INTRAVENOUS SOLUTION 7 %
- *amphotericin b intravenous solution reconstituted 50 mg*
- *aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg*
- AZASAN ORAL TABLET 100 MG, 75 MG
- *azathioprine oral tablet 50 mg*
- *budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml*
- *calcitonin (salmon) nasal solution 200 unit/act*
- *caspofungin acetate intravenous solution reconstituted 50 mg, 70 mg*
- *chlorpromazine hcl oral tablet 10 mg, 25 mg*
- *cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg*
- CLINISOL SF INTRAVENOUS SOLUTION 15 %
- *colistimethate sodium (cba) injection solution reconstituted 150 mg*
- *cromolyn sodium inhalation nebulization solution 20 mg/2ml*
- *cyclophosphamide oral capsule 25 mg, 50 mg*
- *cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg*
- *cyclosporine modified oral solution 100 mg/ml*
- *cyclosporine oral capsule 100 mg, 25 mg*
- *dextrose intravenous solution 10 %, 5 %*
- *diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml*
- ELIGARD SUBCUTANEOUS KIT 22.5 MG, 30 MG, 45 MG, 7.5 MG
- ENGERIX-B INJECTION SUSPENSION 10 MCG/0.5ML, 20 MCG/ML
- ENVARUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG
- *everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg*
- FIRMAGON (240 MG DOSE) SUBCUTANEOUS SOLUTION RECONSTITUTED 120 MG/VIAL
- FIRMAGON SUBCUTANEOUS SOLUTION RECONSTITUTED 80 MG
- GENGRAF ORAL CAPSULE 100 MG, 25 MG
- GENGRAF ORAL SOLUTION 100 MG/ML
- *granisetron hcl oral tablet 1 mg*
- HEPATAMINE INTRAVENOUS SOLUTION 8 %
- IMOVAX RABIES INTRAMUSCULAR INJECTABLE 2.5 UNIT/ML
- INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %
- INTRON A INJECTION SOLUTION 10000000 UNIT/ML, 6000000 UNIT/ML
- INTRON A INJECTION SOLUTION RECONSTITUTED 10000000 UNIT, 18000000 UNIT, 50000000 UNIT

- *ipratropium bromide inhalation solution 0.02 %*
- *ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml*
- ISOLYTE-P IN D5W INTRAVENOUS SOLUTION
- *levolbuterol hcl inhalation nebulization solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25 mg/0.5ml, 1.25 mg/3ml*
- *levocarnitine oral solution 1 gm/10ml*
- *levocarnitine oral tablet 330 mg*
- *methotrexate oral tablet 2.5 mg*
- *methotrexate sodium (pf) injection solution 50 mg/2ml*
- *methotrexate sodium injection solution 50 mg/2ml*
- *mycophenolate mofetil oral capsule 250 mg*
- *mycophenolate mofetil oral suspension reconstituted 200 mg/ml*
- *mycophenolate mofetil oral tablet 500 mg*
- *mycophenolate sodium oral tablet delayed release 180 mg, 360 mg*
- NEPHRAMINE INTRAVENOUS SOLUTION 5.4 %
- NUTRILIPID INTRAVENOUS EMULSION 20 %
- *ondansetron hcl oral solution 4 mg/5ml*
- *ondansetron hcl oral tablet 24 mg, 4 mg, 8 mg*
- *ondansetron oral tablet dispersible 4 mg, 8 mg*
- *pentamidine isethionate inhalation solution reconstituted 300 mg*
- PLENAMINE INTRAVENOUS SOLUTION 15 %
- PREMASOL INTRAVENOUS SOLUTION 10 %
- PROCALAMINE INTRAVENOUS SOLUTION 3 %
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PROSOL INTRAVENOUS SOLUTION 20 %
- PULMOZYME INHALATION SOLUTION 1 MG/ML
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 10 MCG/ML (1ML SYRINGE), 40 MCG/ML, 5 MCG/0.5ML
- SANDIMMUNE ORAL SOLUTION 100 MG/ML
- *sirolimus oral solution 1 mg/ml*
- *sirolimus oral tablet 0.5 mg, 1 mg, 2 mg*
- *tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg*
- TDVAX INTRAMUSCULAR SUSPENSION 2-2 LF/0.5ML
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU
- *tigecycline intravenous solution reconstituted 50 mg*
- *tobramycin inhalation nebulization solution 300 mg/5ml*
- TPN ELECTROLYTES INTRAVENOUS CONCENTRATE
- TRAVASOL INTRAVENOUS SOLUTION 10 %
- TRELSTAR MIXJECT INTRAMUSCULAR SUSPENSION RECONSTITUTED 11.25 MG, 22.5 MG, 3.75 MG
- TREXALL ORAL TABLET 10 MG, 15 MG, 5 MG, 7.5 MG
- TROPHAMINE INTRAVENOUS SOLUTION 10 %
- VARUBI (180 MG DOSE) ORAL TABLET THERAPY PACK 2 X 90 MG
- XATMEP ORAL SOLUTION 2.5 MG/ML
- ZORTRESS ORAL TABLET 1 MG



Local.
Reliable.
Accessible.

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Index

A	
ABELCET INTRAVENOUS SUSPENSION 5 MG/ML.....	201
abiraterone acetate	1
acetylcysteine inhalation solution 10 %, 20 %	201
ACTIMMUNE.....	2
acyclovir sodium intravenous solution 50 mg/ml	201
ADEMPAS	3
AFINITOR DISPERZ.....	4
AFINITOR ORAL TABLET 10 MG	54
albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml ..	201
ALECENSA.....	5
ALKINDI SPRINKLE.....	6
ALUNBRIG.....	8
ALYQ	9
AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG	201
ambrisentan.....	10
AMINOSYN-PF INTRAVENOUS SOLUTION 7 %	201
amphetamine-dextroamphet er.....	11
amphotericin b intravenous solution reconstituted 50 mg.....	201
aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg	201
ARCALYST	12
ARIKAYCE.....	13
armodafinil.....	28
AURYXIA	14
AUSTEDO.....	15
AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT.....	112
AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT.....	112
AYVAKIT	16
AZASAN ORAL TABLET 100 MG, 75 MG	201
azathioprine oral tablet 50 mg.....	201
B	
BALVERSA	17
BETASERON SUBCUTANEOUS KIT	112
bosentan	18
BOSULIF.....	19
BRAFTOVI ORAL CAPSULE 75 MG ...	20
BRUKINSA	21
budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml	201
C	
CABLIVI	22
CABOMETYX	23
calcitonin (salmon) nasal solution 200 unit/act.....	201
CALQUENCE	24
CAPRELSA	25
casprofungin acetate intravenous solution reconstituted 50 mg, 70 mg.....	201
CAYSTON.....	26
chlorpromazine hcl oral tablet 10 mg, 25 mg	201
chlorzoxazone oral tablet 375 mg, 500 mg, 750 mg	73
cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg	201
CINRYZE	27
CLINISOL SF INTRAVENOUS SOLUTION 15 %.....	201
CLOVIQUE	172
colistimethate sodium (cba) injection solution reconstituted 150 mg.....	201
COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG	29
COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG	29
COMETRIQ (60 MG DAILY DOSE).....	29

COPAXONE SUBCUTANEOUS SOLUTION PREFILLED SYRINGE ..	66	ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE	45
COPIKTRA.....	30	ENBREL SUBCUTANEOUS SOLUTION RECONSTITUTED.....	45
CORLANOR ORAL TABLET	31	ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR	45
COSENTYX (300 MG DOSE).....	32	ENDARI	46
COSENTYX SENSOREADY (300 MG). 32		ENGERIX-B INJECTION SUSPENSION 10 MCG/0.5ML, 20 MCG/ML	201
COTELLIC	33	ENSPRYNG	47
cromolyn sodium inhalation nebulization solution 20 mg/2ml	201	ENTRESTO	48
cyclobenzaprine hcl oral	73	ENVARBUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG	201
cyclophosphamide oral capsule 25 mg, 50 mg	201	EPIDIOLEX.....	49
cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg	201	ERLEADA	51
cyclosporine modified oral solution 100 mg/ml	201	erlotinib hcl	52
cyclosporine oral capsule 100 mg, 25 mg	201	ESBRIET	53
CYSTADROPS.....	34	everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg	201
CYSTARAN	35	everolimus oral tablet 2.5 mg, 5 mg, 7.5 mg	54
D		EVRYSDI	55
dalfampridine er	36	F	
DAURISMO	37	FARYDAK	56
deferasirox oral tablet	38	fentanyl citrate buccal lozenge on a handle	57
deferasirox oral tablet soluble.....	38	FINTEPLA.....	58
dextroamphetamine sulfate er	11	FIRAZYR	59
dextroamphetamine sulfate oral.....	11	FIRDAPSE.....	60
dextrose intravenous solution 10 %, 5 %	201	FIRMAGON (240 MG DOSE) SUBCUTANEOUS SOLUTION RECONSTITUTED 120 MG/VIAL... 201	
DIACOMIT.....	39	FIRMAGON SUBCUTANEOUS SOLUTION RECONSTITUTED 80 MG	201
diclofenac sodium external gel 3 %.....	40	G	
dimethyl fumarate oral.....	41	GALAFOLD	61
dimethyl fumarate starter pack.....	41	GAMMAGARD	89
diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml.....	201	GAMMAGARD S/D LESS IGA.....	89
DOJOLVI.....	42	GAMUNEX-C	89
dronabinol	43	GATTEX.....	62
DUPIXENT.....	44		
E			
ELIGARD SUBCUTANEOUS KIT 22.5 MG, 30 MG, 45 MG, 7.5 MG.....	201		
ENBREL MINI.....	45		
ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML	45		

GAVRETO	63	INREBIC.....	83
GENGRAF ORAL CAPSULE 100 MG, 25 MG	201	INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %	201
GENGRAF ORAL SOLUTION 100 MG/ML	201	INTRAROSA.....	84
GILENYA ORAL CAPSULE 0.5 MG....	64	INTRON A INJECTION SOLUTION 10000000 UNIT/ML, 6000000 UNIT/ML	201
GILOTRIF	65	INTRON A INJECTION SOLUTION RECONSTITUTED 10000000 UNIT, 18000000 UNIT, 50000000 UNIT	201
glatiramer acetate	66	ipratropium bromide inhalation solution 0.02 %	202
GOCOVRI	67	ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml	202
granisetron hcl oral tablet 1 mg	201	IRESSA.....	85
H		ISOLYTE-P IN D5W INTRAVENOUS SOLUTION.....	202
HEMADY	70	ISTURISA.....	86
HEPATAMINE INTRAVENOUS SOLUTION 8 %	201	itraconazole oral.....	87, 88
HETLIOZ.....	72	J	
HUMIRA PEDIATRIC CROHNS START SUBCUTANEOUS PREFILLED SYRINGE KIT 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML	74, 75	JAKAFI.....	90
HUMIRA PEN SUBCUTANEOUS PEN- INJECTOR KIT	74, 75	JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG	91
HUMIRA PEN-CD/UC/HS STARTER ..	74, 75	K	
HUMIRA PEN-PS/UV/ADOL HS START SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML	74, 75	KALYDECO.....	92
HUMIRA PEN-PSOR/UEVIT STARTER	74, 75	KESIMPTA.....	93
HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML	74, 75	KISQALI (200 MG DOSE)	94
I		KISQALI (400 MG DOSE)	94
IBRANCE.....	76	KISQALI (600 MG DOSE)	94
ICLUSIG.....	77	KISQALI FEMARA (400 MG DOSE)	95
IDHIFA.....	78	KISQALI FEMARA (600 MG DOSE)	95
IMBRUVICA.....	79	KISQALI FEMARA(200 MG DOSE)	95
IMOVAX RABIES INTRAMUSCULAR INJECTABLE 2.5 UNIT/ML	201	KORLYM	96
INBRIJA	80	KOSELUGO.....	97
INCRELEX.....	81	KUVAN	98
INQOVI	82	L	
		LENVIMA (10 MG DAILY DOSE)	99
		LENVIMA (12 MG DAILY DOSE)	99
		LENVIMA (14 MG DAILY DOSE)	99
		LENVIMA (18 MG DAILY DOSE)	99
		LENVIMA (20 MG DAILY DOSE)	99
		LENVIMA (24 MG DAILY DOSE)	99

LENVIMA (4 MG DAILY DOSE)	99	mycophenolate mofetil oral tablet 500 mg	
LENVIMA (8 MG DAILY DOSE)	99	202
LEUKINE INJECTION SOLUTION		mycophenolate sodium oral tablet delayed	
RECONSTITUTED.....	100	release 180 mg, 360 mg	202
leuprolide acetate injection	101	N	
levabuterol hcl inhalation nebulization		NATPARA.....	113
solution 0.31 mg/3ml, 0.63 mg/3ml, 1.25		NEPHRAMINE INTRAVENOUS	
mg/0.5ml, 1.25 mg/3ml.....	202	SOLUTION 5.4 %	202
levocarnitine oral solution 1 gm/10ml	202	NERLYNX	114
levocarnitine oral tablet 330 mg	202	NEXAVAR.....	115
lidocaine external patch 5 %.....	102	NINLARO.....	116
LONSURF	103	NORTHERA.....	117
LORBRENA.....	104	NUBEQA.....	118
LUPRON DEPOT (1-MONTH).....	101	NUCALA.....	119
LUPRON DEPOT (3-MONTH).....	101	NUEDEXTA.....	120
LUPRON DEPOT (4-MONTH).....	101	NUPLAZID ORAL CAPSULE.....	121
LUPRON DEPOT (6-MONTH).....	101	NUPLAZID ORAL TABLET 10 MG....	121
LYNPARZA ORAL TABLET	105, 106	NUTRILIPID INTRAVENOUS	
M		EMULSION 20 %.....	202
MAVYRET.....	71	O	
MAYZENT	107	OCTAGAM INTRAVENOUS SOLUTION	
MEKINIST	108	1 GM/20ML, 2 GM/20ML	89
MEKTOVI.....	109	octreotide acetate injection solution 100	
methocarbamol oral	73	mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50	
methotrexate oral tablet 2.5 mg	202	mcg/ml, 500 mcg/ml	122
methotrexate sodium (pf) injection solution		ODOMZO	123
50 mg/2ml	202	OFEV	124
methotrexate sodium injection solution 50		OMNITROPE SUBCUTANEOUS	
mg/2ml	202	SOLUTION CARTRIDGE.....	68, 69
methylphenidate hcl er.....	110	OMNITROPE SUBCUTANEOUS	
methylphenidate hcl er (cd)	110	SOLUTION RECONSTITUTED... ..	68, 69
methylphenidate hcl er (la)	110	ondansetron hcl oral solution 4 mg/5ml..	202
methylphenidate hcl er (xr).....	110	ondansetron hcl oral tablet 24 mg, 4 mg, 8	
methylphenidate hcl oral solution.....	110	mg	202
methylphenidate hcl oral tablet.....	110	ondansetron oral tablet dispersible 4 mg, 8	
miglustat.....	111	mg	202
modafinil	28	ONUREG.....	125
mycophenolate mofetil oral capsule 250 mg		OPSUMIT	126
.....	202	ORGOVYX.....	127
mycophenolate mofetil oral suspension		ORLISSA.....	128
reconstituted 200 mg/ml	202	ORKAMBI.....	129
		orphenadrine citrate er	73

OSPHENA 130
oxandrolone oral 131

P

PEGASYS SUBCUTANEOUS SOLUTION
..... 132

PEMAZYRE 133

pentamidine isethionate inhalation solution
reconstituted 300 mg 202

PIQRAY (200 MG DAILY DOSE)..... 134

PIQRAY (250 MG DAILY DOSE)..... 134

PIQRAY (300 MG DAILY DOSE)..... 134

PLENAMINE INTRAVENOUS
SOLUTION 15 % 202

POMALYST 135

PREMASOL INTRAVENOUS SOLUTION
10 % 202

PRIVIGEN INTRAVENOUS SOLUTION
20 GM/200ML 89

PROCALAMINE INTRAVENOUS
SOLUTION 3 % 202

PROGRAF ORAL PACKET 0.2 MG, 1 MG
..... 202

PROLASTIN-C INTRAVENOUS
SOLUTION RECONSTITUTED..... 7

PROMACTA 136

PROSOL INTRAVENOUS SOLUTION 20
% 202

PULMOZYME INHALATION
SOLUTION 1 MG/ML 202

Q

QINLOCK..... 137

R

RABAVERT INTRAMUSCULAR
SUSPENSION RECONSTITUTED... 202

RECOMBIVAX HB INJECTION
SUSPENSION 10 MCG/ML, 10
MCG/ML (1ML SYRINGE), 40
MCG/ML, 5 MCG/0.5ML 202

REGANEX..... 138

REPATHA 139

REPATHA PUSHTRONEX SYSTEM.. 139

REPATHA SURECLICK..... 139

RETACRIT INJECTION SOLUTION

10000 UNIT/ML, 10000
UNIT/ML(1ML), 2000 UNIT/ML, 20000
UNIT/ML, 3000 UNIT/ML, 4000
UNIT/ML, 40000 UNIT/ML 50

RETEVMO 140

REVLIMID 141

ROZLYTREK..... 142

RUBRACA 143

RYDAPT..... 144

S

SAMSCA 145

SANDIMMUNE ORAL SOLUTION 100
MG/ML 202

sapropterin dihydrochloride 146

SIGNIFOR 147

sildenafil citrate oral tablet 20 mg 148

sirolimus oral solution 1 mg/ml 202

sirolimus oral tablet 0.5 mg, 1 mg, 2 mg 202

sofosbuvir-velpatasvir 71

SOMATULINE DEPOT..... 149

SOMAVERT..... 150

SPRYCEL 151

STELARA SUBCUTANEOUS
SOLUTION 45 MG/0.5ML 152

STELARA SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE 152

STIVARGA 153

SUNOSI 154

SUTENT 155

SYMDEKO..... 156

SYMLINPEN 120 SUBCUTANEOUS
SOLUTION PEN-INJECTOR..... 157

SYMLINPEN 60 SUBCUTANEOUS
SOLUTION PEN-INJECTOR..... 157

T

TABRECTA 158
tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg
..... 202

TAFINLAR..... 159

TAGRISO 160

TAKHZYRO..... 161

TALZENNA	162	VENCLEXTA.....	178
TARGRETIN EXTERNAL.....	163	VENCLEXTA STARTING PACK	178
TASIGNA.....	164	VERZENIO.....	179, 180
TAZVERIK.....	165	VITRAKVI.....	181
TDVAX INTRAMUSCULAR		VIZIMPRO	182
SUSPENSION 2-2 LF/0.5ML	202	voriconazole oral.....	183
TECFIDERA.....	166	VOSEVI.....	71
TEGSEDI.....	167	VOTRIENT.....	184
TENIVAC INTRAMUSCULAR		VYNDAMAX.....	185
INJECTABLE 5-2 LFU.....	202	VYVANSE	11
teriparatide (recombinant).....	168	X	
tetrabenazine	169	XALKORI.....	186
THALOMID	170	XATMEP ORAL SOLUTION 2.5 MG/ML	
TIBSOVO	171	202
tigecycline intravenous solution		XELJANZ ORAL TABLET.....	187
reconstituted 50 mg.....	202	XELJANZ XR	187
tobramycin inhalation nebulization solution		XGEVA.....	188
300 mg/5ml.....	202	XOLAIR	189
tolvaptan.....	145	XOSPATA	190
TPN ELECTROLYTES INTRAVENOUS		XPOVIO (100 MG ONCE WEEKLY)...	191
CONCENTRATE	202	XPOVIO (40 MG ONCE WEEKLY)....	191
TRAVASOL INTRAVENOUS SOLUTION		XPOVIO (40 MG TWICE WEEKLY)...	191
10 %	202	XPOVIO (60 MG ONCE WEEKLY)....	191
TRELSTAR MIXJECT		XPOVIO (60 MG TWICE WEEKLY)...	191
INTRAMUSCULAR SUSPENSION		XPOVIO (80 MG ONCE WEEKLY)....	191
RECONSTITUTED 11.25 MG, 22.5 MG,		XPOVIO (80 MG TWICE WEEKLY)...	191
3.75 MG	202	XTANDI ORAL CAPSULE.....	192
TREXALL ORAL TABLET 10 MG, 15		XURIDEN.....	193
MG, 5 MG, 7.5 MG	202	XYREM	194
trientine hcl	172	XYWAV	195
TRIKAFTA.....	173	Y	
TROPHAMINE INTRAVENOUS		YONSA.....	196
SOLUTION 10 %	202	Z	
TUKYSA	174	ZARXIO	197
TURALIO	175	ZEJULA.....	198
TYMLOS	176	ZIEXTENZO	199
U		ZORTRESS ORAL TABLET 1 MG.....	202
UPTRAVI.....	177	ZYKADIA ORAL TABLET	200
V		ZYTIGA ORAL TABLET 500 MG.....	1
VARUBI (180 MG DOSE) ORAL			
TABLET THERAPY PACK 2 X 90 MG			
.....	202		



Local.
Reliable.
Accessible.